

## Immunization Requirements for Preschool – 12<sup>th</sup> Grade

<b>3 Years</b>	<b>4 Years</b>	<b>5 Years</b>	<b>11-15 Years</b>	<b>16 Years and older</b>
<u>DTaP</u> – 4 doses	* <u>DTaP</u> – 4 or 5 doses	* <u>DTaP</u> – 4 or 5 doses	* <u>DTaP</u> – 4 or 5 doses	* <u>DTaP</u> – 4 or 5 doses
Polio ( <u>IPV</u> ) – 3 doses	*Polio ( <u>IPV</u> ) – 3 or 4 doses	*Polio ( <u>IPV</u> ) – 3 or 4 doses	*Polio ( <u>IPV</u> ) – 3 or 4 doses	*Polio ( <u>IPV</u> ) – 3 or 4 doses
<u>MMR</u> – 1 dose	<u>MMR</u> – 2 doses	<u>MMR</u> – 2 doses	<u>MMR</u> – 2 doses	<u>MMR</u> – 2 doses
* <u>Hib</u> – 3 or 4 doses	* <u>Hib</u> – 3 or 4 doses	* <u>Hib</u> – no requirement	<u>Tdap</u> – 1 dose	<u>Tdap</u> – 1 dose
* <u>Hep A</u> - 2 doses effective for the school year beginning on or after <b>July 1, 2018</b>	* <u>Hep A</u> - 2 doses effective for the school year beginning on or after <b>July 1, 2018</b>	* <u>Hep A</u> - 2 doses effective for the school year beginning on or after <b>July 1, 2018</b>	* <u>Hep A</u> -2 doses effective for the school year beginning on or after <b>July 1, 2018</b>	* <u>Hep A</u> 2 doses effective for the school year beginning on or after <b>July 1, 2018</b>
<u>Hep B</u> – 3 doses	<u>Hep B</u> – 3 doses	<u>Hep B</u> – 3 doses	<u>Hep B</u> – 3 doses	<u>Hep B</u> – 3 doses
* <u>Varicella</u> – 1 dose	* <u>Varicella</u> – 2 doses	* <u>Varicella</u> – 2 doses	* <u>Varicella</u> – 2 doses	* <u>Varicella</u> – 2 doses
* <u>Pneumococcal</u> ( <u>PCV</u> ) – 4 doses	* <u>Pneumococcal</u> ( <u>PCV</u> ) – 4 doses	* <u>Pneumococcal</u> ( <u>PCV</u> ) – no requirement	<u>Meningococcal ACWY</u> ( <u>MCV4</u> ) – 1 dose (at 11 years of age)	* <u>Meningococcal ACWY</u> ( <u>MCV4</u> ) – 2 doses for 16 yrs and older effective for the school year beginning on or after <b>July 1, 2018</b>

\*DTaP - Fifth dose not needed if dose #4 given after 4<sup>th</sup> B-day and 6 months from previous dose

\* Polio - Fourth dose not needed if dose #3 given after 4<sup>th</sup> B-day and 6 months from previous dose

\* Hib - last dose must be on or after 12 months of age-not required at age 5 years and older; number of total doses (3 or 4) depends on brand used

\* PCV - last dose must be on or after 12 months of age-not required at age 5 years and older

\* Varicella - no doses needed if history of Chickenpox or Zoster disease is verified by a medical provider

\* Meningococcal ACYW – if initial dose given at 16 yrs or older, will only need one dose (previously known as MCV4)

